

Tobramycin Ophthalmic - Main

Tobramycin Ophthalmic - Drops (Cats)

Tobramycin Ophthalmic - Drops (Dogs)

Tobramycin Ophthalmic - Ointment (Cats)

Tobramycin Ophthalmic - Ointment (Dogs)

GENERAL DESCRIPTION: This drug is an eye medication that is used in dogs and cats primarily to treat bacterial infections. This medication is available in the form of drops as well as an ointment.

WHAT IS THIS DRUG?

An aminoglycoside antibiotic

REASONS FOR PRESCRIBING:

To treat or prevent bacterial infections of the eye, conjunctiva and eyelids

WHAT DOGS/CATS SHOULD NOT TAKE THIS MEDICATION?

Animals with a known hypersensitivity or allergy to this drug should not take this medication

DIRECTIONS:

Read and follow the prescription label carefully.

Give the exact amount prescribed and only as often as directed.

Give this medication for as long as your veterinarian directs.

Do not skip doses or stop giving the medication without consulting your veterinarian.

Dose adjustments may be made based upon an assessment of how your dog or cat is responding clinically.

If your pet requires more than one medication in the same eye, administer the medications 5 minutes apart.

If your pet requires more than one medication, discuss with your veterinarian if it is preferable to administer drops before ointments.

If you are having difficulty applying the medication, contact your veterinarian.

This medication will take effect quickly, in about 1 to 2 hours, but its effects may not be visibly noticeable for a few days.

Call ahead for refills.

Ideally, give this medication at the same time(s) daily.

ADMINISTRATION OF EYE DROPS

Wash your hands before applying any eye medication

Check label directions and shake bottle if necessary

Remove excess discharge from around the eye with a moistened cloth

Remove the cap and avoid contaminating the tip of the medication bottle with the environment or your fingers

Tilt your pet's head back so that they are looking up towards the ceiling

Gently hold the eyelids open and with the other hand holding the medication bottle, rest your hand above the eye

Be sure that the tip of the medication bottle is not pointed towards the eye in case of any sudden movements

Ensure that the tip of the medication bottle does not touch your pet's eye, eyelids, or lashes

Apply the correct number of drops into the eye

Continue to hold your pet's head up and allow them to blink to distribute the drops over the entire eye

Place the cap back on the medication bottle and avoid contaminating the inside of the cap with your fingers

Wash your hands after applying any eye medication

Always praise and reward your pet to encourage cooperation

ADMINISTRATION OF EYE OINTMENTS:

Wash hands prior before applying any eye medication

Check label directions

Remove excess discharge from around the eye with a moistened cloth

Remove the cap and avoid contaminating the tip of the medication tube with the environment or your fingers

Tilt your pet's head back so that they are looking up towards the ceiling

Gently roll down the lower eyelid of the affected eye and while holding the medication tube, rest your hand above the eye

Be sure that the tip of the medication tube is not pointed towards the eye in case of any sudden movements

Ensure that the tip of the medication tube does not touch your pet's eye, eyelids, or lashes

Holding the tube parallel to the eyelid, apply the correct amount of ointment along the inner edge of the lower lid

Gently hold the eyelids shut and massage to disperse the medication

Place the cap back on the medication tube and avoid contaminating the inside of the cap with your fingers

Wash your hands after applying any eye medication

Always praise and reward your pet to encourage cooperation

WHAT IF A DOSE IS MISSED?

If a dose is missed, give it as soon as you can. If it is time already for the next dose, skip the missed dose and go back to the normal schedule. Do not give two doses at the same time.

WHAT TO TELL/ASK YOUR VETERINARIAN BEFORE GIVING THIS MEDICATION?

Talk to your veterinarian about:

When your pet will need to be rechecked

What tests may need to be performed prior to and during treatment with this drug

What are the risks and benefits of using this drug

Tell your veterinarian about:

If your pet has experienced previous eye problems

If your pet has experienced side effects on other drugs/products

If your pet has experienced digestive upset now or ever

If your pet has experienced liver or kidney disease now or ever

If your pet has experienced any other medical problems or allergies now or ever

All medicines and supplements that you are giving your pet or plan to give your pet, including those you can get without a prescription. Your veterinarian may want to check that all of your pet's medications can be given together.

If your pet is pregnant, nursing or if you plan to breed your pet

STORAGE AND WARNINGS:

Tobramycin Ophthalmic should be stored in a childproof container in a cool, dry place at room temperature away from heat and direct sunlight.

Keep this and all medication out of reach of children and pets.

Call your physician immediately if you accidentally take this product.

POTENTIAL SIDE EFFECTS:

This medication may cause the eye to sting for a short period after application

This medication can cause hypersensitivity reactions resulting in marked redness and swelling of the tissues surrounding the eye

This medication can cause the pupil to dilate resulting in sensitivity to light

Prolonged use of any antibiotic may lead to the development of bacterial resistance and overgrowth of other organisms

If these symptoms persist or you notice anything else unusual, contact your veterinarian

This short-acting medication should stop working within 24 hours, although effects may last longer in pets with liver or kidney disease

CAN THIS DRUG BE GIVEN WITH OTHER MEDICATIONS?

Do not give new food or medications without first talking to your veterinarian

If your pet experiences any unusual reactions when taking multiple medications, contact your veterinarian

WHAT SHOULD I DO IF I KNOW OF OR SUSPECT THERE HAS BEEN AN OVERDOSE?

Contact your veterinarian immediately if you pet has accidentally ingested this medication or has received more than the prescribed amount.

WHAT ELSE SHOULD I KNOW?

Notify your veterinarian if your animal's condition does not improve or worsens despite this treatment.

As with all prescribed medicines, Tobramycin Ophthalmic should only be given to the pet for which it was prescribed. It should be given only for the condition for which it was prescribed. Never use skin or ear medications in the eye.

This is just a summary of information about Tobramycin Ophthalmic. If you have any questions or concerns about Tobramycin Ophthalmic or the condition for which it was prescribed, please contact your veterinarian.